

z -CLASSES IN FINITE GROUPS OF CONJUGATE TYPE $(n, 1)$

SHIVAM ARORA AND KRISHNENDU GONGOPADHYAY

ABSTRACT. Two elements in a group G are said to *z -equivalent* or to be in the same *z -class* if their centralizers are conjugate in G . In [KKJ16], it was proved that a non-abelian p -group G can have at most $\frac{p^k-1}{p-1} + 1$ number of z -classes, where $|G/Z(G)| = p^k$. In this note, we characterize the p -groups of conjugate type $(n, 1)$ attaining this maximal number. As a corollary, we characterize p -groups having prime order commutator subgroup and maximal number of z -classes.

1. INTRODUCTION

Two elements in a group G are said to *z -equivalent* or to be in the same *z -class* if their centralizers are conjugate in G . The z -equivalence is an equivalence relation in G and give a partition of G into disjoint subsets. It is coarser than the conjugacy relation. An infinite group may have infinitely many conjugacy classes, but often the number of z -classes is finite and this finiteness of z -classes give a rough idea about dynamical types in a homogeneous space on which G acts, see [Kul07]. There have been some recent work where z -classes in various groups have been investigated, see [Gou06, Kul08, Sin08, GK09, Gon13, GK16]. We note here that a closely related problem is to compute the *genus number* that has been investigated in [Bos13]. Jadhav and Kitture has classified z -classes in p -groups of order up to p^5 , [JK14].

In a recent work Jadhav, Kitture and Kulkarni [KKJ16] has investigated z -classes in finite p -groups. They proved that a non-abelian p -group has at least $p + 2$ z -classes and characterized the groups that attains this lower bound. They also proved that a non-abelian p -group can have at most $\frac{p^k-1}{p-1} + 1$ number of z -classes, where $|G/Z(G)| = p^k$. They proved the following theorem that gives necessary conditions for a p -group G to attain this maximal number of z -classes.

Theorem JKK. [KKJ16] *Let G be a non-abelian group with $|G/Z(G)| = p^k$. If the number of z -classes in G is $\frac{p^k-1}{p-1} + 1$ then either $G/Z(G) \cong C_p \times C_p$ or the following holds:*

- (1) G has no abelian subgroup of index p .
- (2) $G/Z(G)$ is an elementary abelian p -group.

2010 *Mathematics Subject Classification.* 20D15 (Primary); 20E45 (Secondary).

Key words and phrases. conjugacy classes of centralizers, z -classes, p -groups, extraspecial groups.

The authors further gave a counterexample to show that conditions (1) and (2) are not sufficient to attain this bound. It would be interesting to obtain sufficient conditions for a group G to attain the maximal number of z -classes and to characterize such groups.

In this note, we characterize groups of type $(n, 1)$ attaining this bound. To state our result, we recall the following definition by Ito [Ito53].

Definition. [Ito53] *Let G be a finite group. Let $n_1, n_2, \dots, n_r$, where $n_1 > n_2 > \dots > n_r = 1$, be all the numbers which are the indices of the centralizers of elements of G in G . The vector $(n_1, n_2, \dots, n_r)$ is called conjugate type vector of G . A group with the conjugate type vector $(n_1, n_2, \dots, n_r)$ is called a group of type $(n_1, n_2, \dots, n_r)$.*

In [Ito53], Ito characterizes the groups of type $(n, 1)$. Any such group is nilpotent and $n = p^a$ for some prime p . A group of type $(p^a, 1)$ is the direct product of a p -group of the same type and of an abelian group. Thus, the study of groups of type $(n, 1)$ can be reduced to that of p -groups of type $(p^a, 1)$. Ito showed that such a group G contains an abelian normal subgroup A such that G/A is of exponent p .

Here we prove the following.

Theorem 1.1. *Let G be a non-abelian group of type $(n, 1)$ and $[G : Z(G)] = p^k$. Then G has $\frac{p^k - 1}{p - 1} + 1$ z -classes if and only if*

- (1) $G/Z(G)$ is elementary abelian, and
- (2) for all $x \in G \setminus Z(G)$, $Z(C_G(x)) = \langle x, Z(G) \rangle$.

As an application of the above result we have the following.

Corollary 1.2. *Let G be a non-abelian group with $|G/Z(G)| = p^k$, ($k \geq 2$) and $|G'| = p$. Then the number of z -classes in G is $\frac{p^k - 1}{p - 1} + 1$ if and only if G is isoclinic to an extraspecial p -group.*

We prove these results in Section 3.

We note that Corollary 1.2 may be compared with Theorem 5.3.1 of Kitture [Kit14, Section 5.3]. We thank Kitture for letting us know about his result and many comments. We are also thankful to Silvio Dolfi for useful comments and suggestions on a first draft of this note.

2. PRELIMINARIES

We recall here a few basic facts from Kulkarni [Kul07] that will be used in the proof.

Let G be a group. We denote $C_G(x)$ to be the centralizer of an element x in G .

Definition 2.1. *Let G be a group. Two elements $x, y \in G$ are said to be z -equivalent if their centralizers are conjugate, i.e. there exist $g \in G$ such that,*

$$g^{-1}C_G(x)g = C_G(y).$$

Let G be a group acting on a set X . For $x \in X$ let $G(x)$ denote the orbit of x and G_x be the stabilizer subgroup of x :

$$G(x) = \{y \in X : \text{there exists } g \in G \text{ and } y = gx\}.$$

$$G_x = \{g \in G : gx = x\}.$$

We have a partition of X as:

$$X = \bigcup_{x \in X} G(x).$$

Let $y \in G(x)$ be some element, then for some $g \in G$ we have $y = gx$ and $G_y = gG(x)g^{-1}$.

For subgroups A and B in G write $A \sim_g B$ if they are conjugate. This gives us the following.

Definition 2.2. Let $x, y \in X$ be any two element, they are said to be in same orbit class if $G_x \sim_g G_y$. Denote this equivalence relation as $x \sim_0 y$ and denote $R(x)$ as the equivalence class of x .

This gives another partition of X :

$$X = \bigcup_{G_x \sim_g G_y} R(x) = \bigcup_{G_x = G_y} R(x).$$

The second inequality is easy to note, since if $x \sim_0 y$ then for some $z \in X$, we have $C_G(x) = C_G(z)$.

Let F_x denotes the points fixed by G_x i.e.

$$F_x = \{y \in X : G_y \supset G_x\}.$$

Let F'_x denotes the set $F'_x = \{y \in X : G_y = G_x\}$. Define $W_x = N_x/G_x$, where N_x is the normalizer of the G_x in G . Then we have the following theorem(refer [Kul07]).

Theorem 2.3. Let G act on a set X . Then the map $\phi : G/G_x \times F'_x \rightarrow R(x)$, $\phi(gG_x, y) = gy$ is well defined and induces a bijection, $\bar{\phi} : (G/G_x \times F'_x)/W_x \rightarrow R(x)$.

When we take our space $X = G$, the group itself, then $R(x)$ in above notation becomes the z -classes and the above theorem gives us the following equation, which is used to calculate the size of a z -class: for any $x \in G$ we have

$$|z - \text{class of } x| = [G : N_G(C_G(x))] \cdot |F'_x|,$$

where $C_G(x)$ is the centralizer of x and $F'_x = \{y \in G : C_G(y) = C_G(x)\}$.

The following theorem states that that number of z -classes is invariant under the isoclinism. For basic notions on isoclinism, see [Hal40, KKJ16].

Theorem 2.4. [KKJ16] Let G_1 and G_2 be isoclinic groups, with an isoclinism (ϕ, ψ) . Then the isoclinism (ϕ, ψ) induces a bijection between the z -classes in G_1 and G_2 .

Finally we note the following theorem by Hall [Hal40] that is used in proof of Theorem 1.2.

Theorem 2.5. Every group is isoclinic to a group whose center is contained in the commutator subgroup.

3. PROOF OF THE MAIN RESULTS

It was proved in [KKJ16, Lemma 3.1] that if G is a finite group with $|G/Z(G)| = p^k$, then G is isoclinic to a finite p -group. Since number of z -classes is invariant under isoclinism, hence it is enough to prove our results assuming that G is a p -group.

Lemma 3.1. *Let G be a non-abelian p -group with $[G : Z(G)] = p^k$. If*

- (1) $G/Z(G)$ is elementary abelian, and
- (2) for all $z \in G \setminus Z(G)$, $Z(C_G(x)) = \langle x, Z(G) \rangle$,

then G has $\frac{p^k-1}{p-1} + 1$ z -classes.

Proof. Let $x \in G \setminus Z(G)$. Since $G/Z(G)$ is abelian, hence $C_G(x) \trianglelefteq G$, i.e. $[G : N(C_G(x))] = 1$. Consider

$$F'_x = \{y \in G : C_G(y) = C_G(x)\}.$$

By definition $F'_x \subseteq Z(C_G(x)) \setminus Z(G) = \langle x, Z(G) \rangle \setminus Z(G)$. Obviously $C_G(x) = C_G(xt)$ for $t \in Z(G)$. Also since exponent of $G/Z(G)$ is p , thus $C_G(x) = C_G(x^i)$ for any $i \in \{1, 2, \dots, p-1\}$. Hence $\langle x, Z(G) \rangle \setminus Z(G) \subseteq F'_x$. Hence

$$F'_x = Z(C_G(x)) \setminus Z(G) = \langle x, Z(G) \rangle \setminus Z(G) = xZ(G) \cup \dots \cup x^{p-1}Z(G).$$

This implies, $|F'_x| = (p-1)|Z(G)|$. Therefore,

$$|z\text{-class of } x| = [G : N_G(C_G(x))] \cdot |F'_x| = 1 \cdot (p-1)|Z(G)|.$$

Thus $G \setminus Z(G)$ contributes to a total $\frac{p^k-1}{p-1}$ number of z -classes of G . Clearly, $Z(G)$ is also a z -class. Hence, G has $\frac{p^k-1}{p-1} + 1$ z -classes. $\square$

Corollary 3.2. *Let G be a non-abelian p -group with $|G/Z(G)| = p^k$. If*

- (1) G has no abelian subgroup of order exceeding $p|Z(G)|$.
- (2) $G/Z(G)$ is an elementary abelian p -group.

Then G has $\frac{p^k-1}{p-1} + 1$ number of z -classes.

Proof. We claim $C_G(x) = \langle x, Z(G) \rangle$, and the rest follow from Lemma 3.1.

Clearly $\langle x, Z(G) \rangle \subseteq C_G(x)$. Let us assume $y \notin \langle x, Z(G) \rangle$ and $y \in C_G(x)$. But then $\langle x, y, Z(G) \rangle$ is an abelian subgroup and $|\langle x, y, Z(G) \rangle| > p|Z(G)|$. Thus $C_G(x) = \langle x, Z(G) \rangle$. $\square$

3.1. Proof of Theorem 1.1.

Proof. Let G be a non-abelian group of type $(n, 1)$ and G has $\frac{p^k-1}{p-1} + 1$ z -classes, where $[G : Z(G)] = p^k$. (1) follows from the Theorem JKK. We prove (2).

Let $x \in G \setminus Z(G)$, then for any $t \in Z(G)$, $C_G(x) = C_G(xt)$. Also for any m relatively prime to order of x , we have

$$C_G(x) = C_G(\langle x \rangle) = C_G(\langle x^m \rangle)$$

Thus z -class of x contains,

$$xZ(G) \cup x^2Z(G) \cup \dots \cup x^{p-1}Z(G),$$

and hence, for all $x \in G \setminus Z(G)$, the size of z class is at least $(p-1)|Z(G)|$. Since $Z(G)$ is also a z -class, hence it follows from the hypothesis that $G \setminus Z(G)$ constitute exactly $\frac{p^k-1}{p-1}$ z -classes. Now, $|G \setminus Z(G)| = |G| - |Z(G)|$. Hence, we must have for $x \in G \setminus Z(G)$,

$$|z - \text{class of } x| = (p-1)|Z(G)|.$$

Now, it is clear that for $x \in G \setminus Z(G)$, $\langle x, Z(G) \rangle \subseteq Z(C_G(x))$. Now let $y \in Z(C_G(x))$ such that $y \notin \langle x, Z(G) \rangle$. Then a simple observation shows that $C_G(y) \supseteq C_G(x)$. But since G is of type $(n, 1)$ we have $|C_G(x)| = |C_G(y)|$, thus we must have $C_G(x) = C_G(y)$. This implies $y \in (z - \text{class of } x)$ and since $y \notin \langle x, Z(G) \rangle$, thus $|z - \text{class of } x| > (p-1)|Z(G)|$, which is a contradiction. Thus our assumption is incorrect and $Z(C_G(x)) = \langle x, Z(G) \rangle$ for all $x \in G \setminus Z(G)$.

The converse part follows from Lemma 3.1. This completes the proof. $\square$

3.2. Proof of Corollary 1.2.

Proof. : Without loss of generality, using Theorem 2.5, we assume up to isoclinism that, G is a group such that $Z(G) \subseteq G'$. Let G has $\frac{p^k-1}{p-1} + 1$ z -classes. Since $|G'| = p$ and in a non-abelian p -group $|Z(G)| > 1$, it follows that $G' = Z(G)$; hence, $|G'| = |Z(G)| = p$. Therefore G and $G/Z(G)$ have the same exponent. Thus G is an extra-special p -group.

Conversely, let G be an extraspecial p -group. So, $|Z(G)| = |G'| = p$. For $x \in G \setminus Z(G)$, define the map:

$$\begin{aligned} \phi_x : G &\rightarrow G' \\ g &\mapsto [x, g] \end{aligned}$$

As $G/Z(G)$ is elementary abelian, ϕ_x is a homomorphism with $\ker(\phi_x) = C_G(x)$ and since $|G'| = p$, the map is also surjective. Hence we get

$$G/C_G(x) \cong G'$$

Thus each $C_G(x)$ is a maximal subgroup of index p in G . This shows that extraspecial p -groups are of type $(n, 1)$.

Now, we claim that $Z(C_G(x)) = \langle x, Z(G) \rangle$. This follows from the fact that $G/Z(G)$ can be equipped with an alternating non-degenerate form induced by the commutator, see the proof of [Cra08, Theorem 3.14] or [Wil09, p. 83]. The claim follows from [Cra08, Lemma 3.11] noting that $Z(C_G(x))/Z(G)$ is the radical of co-dimension one subspace $C_G(x)/Z(G)$.

The result now follows from Theorem 1.1. $\square$

REFERENCES

- [Bos13] Anirban Bose. On the genus number of algebraic groups. *J. Ramanujan Math. Soc.*, 28(4):443–482, 2013.
- [Cra08] David A. Craven. The theory of p -groups. <http://web.mat.bham.ac.uk/D.A.Craven/docs/lectures/pgroups.pdf>, 2008.
- [GK09] Krishnendu Gongopadhyay and Ravi S. Kulkarni. z -classes of isometries of the hyperbolic space. *Conform. Geom. Dyn.*, 13:91–109, 2009.

- [GK16] Krishnendu Gongopadhyay and Ravi S. Kulkarni. The z -classes of isometries. *Journal of Indian Mathematical Society*, 81(3-4), 2016.
- [Gon13] Krishnendu Gongopadhyay. The z -classes of quaternionic hyperbolic isometries. *J. Group Theory*, 16(6):941–964, 2013.
- [Gou06] Rony Gouraige. *Z-classes in central simple algebras*. ProQuest LLC, Ann Arbor, MI, 2006. Thesis (Ph.D.)—City University of New York.
- [Hal40] P. Hall. The classification of prime-power groups. *Journal für die reine und angewandte Mathematik*, 182:130–141, 1940.
- [Ito53] Noboru Ito. On finite groups with given conjugate types I. *Nagoya Math. J.*, 6:17–28, 1953.
- [JK14] Vikas S. Jadhav and Rahul D. Kitture. z -classes in p -groups less than equal to p^5 . *Bull. Allahabad. Soc.*, 29:179–194, 2014.
- [Kit14] Rahul D. Kitture. *z -Classes in finite p -groups*. 2014. Thesis (Ph.D.)—University of Pune.
- [KKJ16] Ravindra Kulkarni, Rahul Dattatraya Kitture, and Vikas S. Jadhav. z -classes in groups. *Journal of Algebra and Its Applications*, 15(6):165013 (16 pages), 2016.
- [Kul07] Ravi S. Kulkarni. Dynamical types and conjugacy classes of centralizers in groups. *J. Ramanujan Math. Soc.*, 22(1):35–56, 2007.
- [Kul08] Ravi S. Kulkarni. Dynamics of linear and affine maps. *Asian J. Math.*, 12(3):321–343, 2008.
- [Sin08] Anupam Singh. Conjugacy classes of centralizers in G_2 . *J. Ramanujan Math. Soc.*, 23(4):327–336, 2008.
- [Wil09] Robert A. Wilson. *The finite simple groups*, volume 251 of *Graduate Texts in Mathematics*. Springer-Verlag London, Ltd., London, 2009.

INDIAN INSTITUTE OF SCIENCE EDUCATION AND RESEARCH (IISER) MOHALI, KNOWLEDGE CITY, , SECTOR 81, S.A.S. NAGAR, PUNJAB 140306, INDIA
E-mail address: shivam.iiserm@gmail.com

INDIAN INSTITUTE OF SCIENCE EDUCATION AND RESEARCH (IISER) MOHALI, KNOWLEDGE CITY, SECTOR 81, S.A.S. NAGAR, PUNJAB 140306, INDIA
E-mail address: krishnendug@gmail.com, krishnendu@iisermohali.ac.in